ESTUARINE COASTAL AND SHELF SCIENCE

Volume 35
1992

ACADEMIC PRESS
London San Diego
New York Boston
Sydney Tokyo Toronto

Published by Academic Press Limited in association with
the Estuarine and Coastal Sciences Association

ISSN 0272–7714

er.
C
6
E78

Copyright © 1992, by Academic Press Limited

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without prior permission from the publishers.

Printed in Great Britain

Contents of Volume 35

Number 1, July 1992

Palacín, C., Gili, J.-M. and Martín, D., Evidence for coincidence of meiofauna spatial heterogeneity with eutrophication processes in a shallow-water Mediterranean bay 1

Wright, L. D., Boon, J. D., Xu, J. P. and Kim, S. C., The Bottom Boundary Layer of the Bay Stem Plains Environment of Lower Chesapeake Bay 17

Millward, G. E., Glegg, G. A. and Morris, A. W., Zn and Cu Removal Kinetics in Estuarine Waters 37

Bach, P., Legendre, P., Amanieu, M. and Lasserre, G., Strategy of eel (*Anguilla anguilla* L.) exploitation in the Thau lagoon. 55

Johns, B., Rao, A. D. and Rao, G. S., On the Occurrence of Upwelling along the East Coast of India 75

Gammelsrød, T., Variation in Shrimp Abundance on the Sofala Bank, Mozambique, and its Relation to the Zambezi River Runoff 91

Notes and Discussions

Shillabeer, N., Hart, B. and Riddle, A. M., The Use of a Mathematical Model to Compare Particle Size Data Derived by Dry-Sieving and Laser Analysis . . . 105

Number 2, August 1992

Shetye, S. R. and Gouveia, A. D., On the Role of Geometry of Cross-Section in Generating Flood-Dominance in Shallow Estuaries 113

Newell, S. Y., Hopkinson, C. S. and Scott, L. A., Patterns of Nitrogenase Activity (Acetylene Reduction) Associated with Standing, Decaying Shoots of *Spartina alterniflora* 127

Takada, H., Ishiwatari, R. and Ogura, N., Distribution of Linear alkylbenzenes (LABs) and Linear alkylbenzene-sulphonates (LAS) in Tokyo Bay Sediments . 141

Garvine, R. W., McCarthy, R. K. and Wong, K.-C., The Axial Salinity Distribution in the Delaware Estuary and its Weak Response to River Discharge 157

Dye, A. H., Recruitment Dynamics and Growth of the Barnacle *Tetraclita Serrata* on the East Coast of Southern Africa 167

Wilber, D. H., Associations Between Freshwater Inflows and Oyster Productivity in Apalachicola Bay, Florida. 179

REVELANTE, N. and GILMARTIN, M., The Lateral Advection of Particulate Organic Matter from the Po Delta Region During Summer Stratification, and its Implications for the Northern Adriatic 191

NOTES AND DISCUSSIONS

NEIRA, F. J. and POTTER, I. C., Movement of Larval Fishes Through the Entrance Channel of a Seasonally Open Estuary in Western Australia 213

NUMBER 3, SEPTEMBER 1992

RENSHUN, Z., Suspended sediment transport processes on tidal mud flat in Jiangsu Province, China 225

LEBO, M. E. and SHARP, J. H., Modeling phosphorus cycling in a well-mixed coastal plain estuary 235

ALLANSON, B. R., SKINNER, D. and IMBERGER, J., Flow in prawn burrows 253

PUŠKARIĆ, S., FOWLER, S. W. and MIQUEL, J.-C., Temporal changes in particulate flux in the northern Adriatic Sea 267

BORSA, P. and MILLET, B., Recruitment of the Clam *Ruditapes decussatus* in the lagoon of Thau, Mediterranean 289

LAW, C. S., REES, A. P. and OWENS, N. J. P., Nitrous Oxide: Estuarine sources and atmospheric flux 301

WEBER, D. E., FLEMER, D. A. and BUNDRICK, C. M., Comparison of the effects of drilling fluid on macrobenthic Invertebrates associated with the Seagrass, *Thalassia testudinum*, in the laboratory and field 315

NUMBER 4, OCTOBER 1992

PILLAY, S., GARDNER, L. R. and KJERFVE, B., The Effect of Cross-Sectional Velocity and Concentration Variations on Suspended Sediment Transport Rates in Tidal Creeks 331

BROCHE, P. and FORGET, P., Has the Influence of Surface Waves on Wind Stress to be Accounted for in Modelling the Coastal Circulation 347

RIDD, P. V., A Sediment Level Sensor for Erosion and Siltation Detection . . . 353

PANDARINATH, K. and NARAYANA, A. C., Clay Minerals and Trace Metal Association in the Gangolli Estuarine Sediments, West Coast of India 363

KERNER, M. and WALLMANN, K., Remobilization Events Involving Cd and Zn from Intertidal Flat Sediments in the Elbe Estuary During the Tidal Cycle 371

HIRCHE, H.-J., Egg Production of *Eurytemora affinis*—Effect of K-Strategy . . . 395

DOLLAR, S. J. and ATKINSON, M. J., Effects of Nutrient Subsidies from Groundwater to Nearshore Marine Ecosystems off the Island of Hawaii 409

De Madariaga, I., González-Azpiri, L., Villate, F. and Orive, E., Plankton Responses to Hydrological Changes Induced by Freshets in a Shallow Mesotidal Estuary 425

Number 5, November 1992

Conlan, K., White, K. N. and Hawkins, S. J., The Hydrography and Ecology of a Re-Developed Brackish-Water Dock 435

Orson, R. A. and Howes, B. L., Salt Marsh Development Studies at Waquoit Bay, Massachusetts: Influence of geomorphology on long-term plant community structure 453

Wolanski, E., Gibbs, R., Ridd, P. and Mehta, A., Settling of Ocean-dumped Dredged Material, Townsville, Australia 473

Llansó, R. J., Effects of Hypoxia on Estuarine Benthos: the Lower Rappahannock River (Chesapeake Bay), a Case Study 491

Althausen, J. D. and Kjerfve, B., Distribution of Suspended Sediment in a Partially Mixed Estuary, Charleston Harbor, South Carolina, U.S.A. 517

Laybourn-Parry, J., Rogerson, A. and Crawford, D. W., Temporal Patterns of Protozooplankton Abundance in the Clyde and Loch Striven 533

Number 6, December 1992

Cyrus, D. P., Blaber, S. J. M., Turbidity and Salinity in a Tropical Northern Australian Estuary and their Influence on Fish Distribution 545

Fonseca, M. S., Cahalan, J. A., A Preliminary Evaluation of Wave Attenuation by Four Species of Seagrass 565

Fichez, R., Jickells, T. D., Edmunds, H. M., Algal Blooms in High Turbidity, a Result of the Conflicting Consequences of Turbulence on Nutrient Cycling in a Shallow Water Estuary 577

Thessalou-Legaki, M., Reproductive Variability of *Parapandalus narval* (Crustacea: Decapoda) Along a Depth Gradient 593

Blaber, S. J. M., Brewer, D. T., Salini, J. P., Kerr, J. D., Conacher, C., Species Composition and Biomasses of Fishes in Tropical Seagrasses at Groote Eylandt, Northern Australia 605

Abreu, P. C., Biddanda, B. B., Odebrecht, C., Bacterial Dynamics of the Patos Lagoon Estuary, Southern Brazil (32°S, 52°W): Relationship with Phytoplankton Production and Suspended Material 621

Freire, J., González-Gurriarán, E., Olaso, I., Spatial Distribution of *Munida intermedia* and *M. sarsi* (Crustacea: Anomura) on the Galician Continental Shelf (NW Spain): Application of Geostatistical Analysis 637

ABREU, P. C., BIDDANDA, B. B. and ODEBRECHT, C., Bacterial dynamics of the Patos Lagoon estuary, southern Brazil (32°S, 52°W): relationship with phytoplankton production and suspended material, 621
Acetylene reduction; saltmarsh; *Spartina alterniflora*; decomposition; clay deposition; nitrogen fixation, 127
Adriatic Sea; eutrophication; phytoplankton; chlorophyll *a* distribution; microdetritus; stratification; advection, 191
Advection; Adriatic Sea; eutrophication; phytoplankton; chlorophyll *a* distribution; microdetritus; stratification, 191
Algal bloom, nutrient cycling; estuary, 577
ALLANSON, B. R., SKINNER, D. and IMBERGER, J., Flow in prawn burrows, 253
ALTHAUSEN, J. D. and KJERFVE, B., Distribution of suspended sediment in a partially mixed estuary, Charleston Harbor, South Carolina, U.S.A., 517
AMANIEU, M. *see* BACH, P., 55
Annual cycle; particulate flux; northern Adriatic Sea, 267
Anthropogenic; nutrients; groundwater; Hawaii; mixing model, 409
Apalachicola Bay; oyster fishery; freshwater inflows, 179
ATKINSON, M. J., *see* DOLLAR, S. J., 409
Atmospheric flux; Tamar estuary; South West England; nitrous oxide; nutrients; nitrification; denitrification; estuarine sediment, 301
Attenuation; habitat; function; seagrass; waves, 565
Australia coast; dredging; sea dumping; settling; sedimentation; mud suspension, 473
Australia coast; tropical seagrass; fish; species composition; biomass; Gulf of Carpentaria, 605
BACH, P., LEGENDRE, P., AMANIEU, M. and LASSERRE, G., Strategy of eel (*Anguilla anguilla* L.) exploitation in the Thau lagoon, 55
Bacteria; phytoplankton; protozoa; seston; particulate organic carbon; estuaries; Brazil coast; South America South coast, 621
Barnacles; intertidal; southern Africa; recruitment; growth, 167
Bass Creek, SC; North Inlet, SC; Suspended sediment concentration; sediment transport rate; tidal estuary; *Spartina* marsh; errors; Long Term Ecological Research (LTER), 331
Bay of Bengal; numerical model; coastal upwelling; estuarial discharge, 75
Bed friction; wave-current interactions; internal waves; estuaries; benthic boundary layer, 17
Benthic boundary layer; bed friction; wave-current interactions; internal waves; estuaries, 17
Bernoulli effect; induced laminar flow; Reynolds number; tube aspect ratio; boundary layer; suspensoids; thalassinidea; burrows, 253
BIDDANDA, B. B., *see* ABREU, P. C., 621
Biomass; Gulf of Carpentaria; Australia coast; tropical seagrass; fish; species composition, 605
Bivalves; *Ruditapes*; Mediterranean; recruitment; larval transport; coastal lagoon, 289
BLABER, S. J. M., BREWER, D. T., SALINI, J. P., KERR, J. D. and CONACHER, C., Species composition and biomasses of fishes in tropical seagrasses at Groote Eylandt, northern Australia, 605
BLABER, S. J. M., *see* CYRUS, D. P., 545
BOON, J. D., *see* WRIGHT, L. D., 17
BORSA, P. and MILLET, B., Recruitment of the clam *Ruditapes decussatus* in the lagoon of Thau, Mediterranean, 289
Boundary layer; suspensoids; thalassinidea; burrows; Bernoulli effect; induced laminar flow; Reynolds number; tube aspect ratio, 253
Brazil coast; South America South coast; bacteria; phytoplankton; protozoa; seston; particulate organic carbon; estuaries, 621
BREWER, D. T., *see* BLABER, S. J. M., 605
BROCHE, P. and FORGET, P., has the influence of surface waves on wind stress to be accounted for in modelling the coastal circulation, 347
BUNDRICK, C. M., *see* WEBER, D. E., 315
Burrows; Bernoulli effect; induced laminar flow; Reynolds number; tube aspect ratio; boundary layer; suspensoids; thalassinidea, 253
Burrowing behaviour; Chesapeake Bay, U.S.A. coast; dissolved oxygen; macrobenthos; population dynamics, 491

Cahalan, J. A., *see* Fonseca, M. S., 565
Calcium; Gangolli estuary; clay minerals; trace elements; organic matter, 363
Chesapeake Bay, U.S.A. coast; dissolved oxygen; macrobenthos; population dynamics; burrowing behaviour, 491
Chlorophyll *a* distribution; microdetritus; stratification; advection; Adriatic Sea; eutrophication; phytoplankton, 191
Ciliates; dinoflagellates; Loch Striven; Clyde; Scotland coast; estuarine; flagellates; ciliates; dinoflagellates; flocs, 533
Clay deposition; nitrogen fixation; acetylene reduction; saltmarsh; *Spartina alterniflora*; decomposition, 127
Clay minerals; trace elements; organic matter; calcium; Gangolli estuary, 363
Clyde; Scotland coast; estuarine; flagellates; ciliates; dinoflagellates; flocs; Loch Striven, 533
Coastal lagoon; bivalves; *Ruditapes*; Mediterranean; recruitment; larval transport, 289
Coastal upwelling; estuarial discharge; Bay of Bengal; numerical model, 75
Community; drilling mud; Santa Rosa Sound, Florida; macrobenthos; seagrass, 315
Conacher, C., *see* Blaber, S. J. M., 605
Conlan, K., White, K. N. and Hawkins, S. J., The hydrography and ecology of a re-developed brackish-water dock, 435
Contingency periodogram; path analysis; eel; fishing gear; fishing effort; fishing strategy; Thau lagoon, 55
Crawford, D. W., *see* Laybourn-Parry, J., 533
Cyrus, D. P., Blaber, S. J. M., Turbidity and salinity in a tropical northern Australian estuary and their influence on fish distribution, 545
Decomposition; clay deposition; nitrogen fixation; acetylene reduction; saltmarsh; *Spartina alterniflora*, 127
Delaware Bay; phosphorus; models; estuaries; mass balance; regeneration, 235
Delaware Bay; salinity variations; estuaries; river flow; diffusivity; effective longitudinal, 157
de Madariaga, I., González-Azpiri, L., Villate, F. and Orive, E., Plankton responses to hydrological changes induced by freshets in a shallow mesotidal estuary, 425
Denitrification; estuarine sediment; atmospheric flux; Tamar estuary; South West England; nitrous oxide; nutrients; nitrification, 301
Density and variations; Mediterranean coast; nematoda; meiobenthos; eutrophication; population, 1
Deposition; removal of contaminants; mudland sandflats; microbial degradation; trace elements; intertidal environment, 371
Depth; Pandalidae; *Parapandalus narval*; Greece coast; fecundity; reproductive effort, 593
Diffusivity; effective longitudinal; Delaware Bay; salinity variations; estuaries; river flow, 157
Dinoflagellates; flocs; Loch Striven; Clyde; Scotland coast; estuarine; flagellates; ciliates, 533
Dissolved oxygen; macrobenthos; population dynamics; burrowing behaviour; Chesapeake Bay, U.S.A. coast, 491
Docks; lagoons; pollution; plankton; England coast; estuaries, 435
Dollar, S. J. and Atkinson, M. J., Effects of nutrient subsidies from groundwater to nearshore marine ecosystems off the island of Hawaii, 409
Dredging; sea dumping; settling; sedimentation; mud suspension; Australia coast, 473
Drilling mud; Santa Rosa Sound, Florida; macrobenthos; seagrass; community, 315
Dye, A. H., Recruitment dynamics and growth of the barnacle *Tetraclita serrata* on the east coast of southern Africa, 167
Edmunds, H. M., *see* Fichez, R., 577
Eel; fishing gear; fishing effort; fishing strategy; Thau lagoon; contingency periodogram; path analysis, 55
Effective longitudinal; Delaware Bay; salinity variations; estuaries; river flow; diffusivity, 157
Egg production; reproductive strategy; *Eurytemora*, 395
Electromagnetics; erosion; siltation; instruments, 353
England coast; estuaries; docks; lagoons; pollution; plankton, 435
England coast; trace metals; kinetics; estuaries, 37
Erosion; siltation; instruments; electromagnetics, 353

Errors; Long Term Ecological Research (LTER); North Inlet, SC; Bass Creek, SC; suspended sediment concentration; sediment transport rate; tidal estuary; *Spartina* marsh, 331
Estuarial discharge; Bay of Bengal; numerical model; coastal upwelling, 75
Estuaries; benthic boundary layer; bed friction; wave-current interactions; internal waves, 17
Estuaries; Brazil coast; South America South coast; bacteria; phytoplankton; protozoa; seston; particulate organic carbon, 621
Estuaries; docks; lagoons; pollution; plankton; England coast, 435
Estuaries; England coast; trace metals; kinetics, 37
Estuaries; fish larvae; transport; Western Australian coast, 213
Estuaries; mass balance; regeneration; Delaware Bay; phosphorus; models, 235
Estuaries; river flow; diffusivity; effective longitudinal; Delaware Bay; salinity variations, 157
Estuarine circulation; tides; models, 113
Estuarine; flagellates; ciliates; dinoflagellates; flocs; Loch Striven; Clyde; Scotland coast, 533
Estuarine habitat; seasonal patterns; turbidity; salinity; temperature; fish distribution, 545
Estuarine sediment; atmospheric flux; Tamar estuary; South West England; nitrous oxide; nutrients; nitrification; denitrification, 301
Estuary; algal bloom; nutrient cycling, 577
Estuary, partially mixed; slack water; suspended sediment; tides; transmissivity; turbidity maximum zone; U.S.A. coast, fresh water discharge, 517
Eurytemora; egg production; reproductive strategy, 395
Eutrophication; phytoplankton; chlorophyll *a* distribution; microdetritus; stratification; advection; Adriatic Sea, 191
Eutrophication; population; density and variations; Mediterranean coast; nematoda; meiobenthos, 1
Fecundity; reproductive effort; depth; Pandalidae; *Parapandalus narval*; Greece coast, 593
Fichez, R., Jickells, T. D. and Edmunds, H. M., Algal blooms in high turbidity, a result of the conflicting consequences of turbulence on nutrient cycling in a shallow water estuary, 577
Fish distribution; estuarine habitat; seasonal patterns; turbidity; salinity; temperature, 545
Fish larvae; transport; Western Australian coast; estuaries, 213
Fish; species composition; biomass; Gulf of Carpentaria; Australia coast; tropical seagrass, 605
Fishing effort; fishing strategy; Thau lagoon; contingency periodogram; path analysis, eel; fishing gear, 55
Fishing gear; fishing effort; fishing strategy; Thau lagoon; contingency periodogram; path analysis; eel, 55
Fishing strategy; Thau lagoon; contingency periodogram; path analysis; eel; fishing gear; fishing effort, 55
Flagellates; ciliates; dinoflagellates; flocs; Loch Striven; Clyde; Scotland coast; estuarine, 533
Flemer, D. A., *see* Weber, D. E., 315
Flocs; Loch Striven; Clyde; Scotland coast; estuarine; flagellates; ciliates; dinoflagellates, 533
Fonseca, M. S. and Cahalan, J. A., A preliminary evaluation of wave attenuation by four species of seagrass, 565
Forget, P., *see* Broche, P., 347
Fowler, S. W., *see* Puškarić, S., 267
Freire, J., González-Gurriarán, E. and Olaso, I., Spatial distribution of *Munida intermedia* and *M. sarsi* (Crustacea: Anomura) on the Galician continental shelf (NW Spain): application of geostatistical analysis, 637
Freshets; natural perturbations; Gernika estuary; plankton dynamics; hydrography, 425
Fresh water discharge; partially mixed estuary; slack water; suspended sediment; tides; transmissivity; turbidity maximum zone; U.S.A. coast, 517
Freshwater inflows; Apalichicola Bay; oyster fishery, 179
Function; seagrass; waves; attenuation; habitat, 565
Galician continental shelf; NW Spain; geostatistical analysis; variograms; kriging; spatial distribution; *Munida intermedia*; *Munida sarsi*, 637
Gammelsrød, T., Variation in shrimp abundance on the Sofala Bank, Mozambique, and its relation to the Zambezi River runoff, 91
Gangolli estuary; clay minerals; trace elements; organic matter; calcium, 363

GARDNER, L. R., *see* PILLAY, S., 331
GARVINE, R. W., MCCARTHY, R. K. and WONG, K.-C., The axial salinity distribution in the Delaware estuary and its weak response to river discharge, 157
Geomorphology; plant community structure; New England Coast; paleoecology, 453
Geostatistical analysis; variograms; kriging; spatial distribution; *Munida intermedia*; *Munida sarsi*; Galician continental shelf; NW Spain, 637
Gernika estuary; plankton dynamics; hydrography; freshets; natural perturbations, 425
GIBBS, R., *see* WOLANSKI, E., 473
GILI, J.-M., *see* PALACÍN, C., 1
GILMARTIN, M., *see* REVELANTE, N., 191
GLEGG, G. A., *see* MILLWARD, G. E., 37
GONZÁLEZ-AZPIRI, L., *see* DE MADARIAGA, I., 425
GONZÁLEZ-GURRIARÁN, E., *see* FREIRE, J., 637
GOUVEIA, A. D., *see* SHETYE, S. R., 113
Greece coast; fecundity; reproductive effort; depth; Pandalidae; *Parapandalus narval*, 593
Groundwater; Hawaii; mixing model; anthropogenic; nutrients, 409
Growth; barnacles; intertidal; southern Africa; recruitment, 167
Gulf of Carpentaria; Australia coast; tropical seagrass; fish; species composition; biomass, 605
Habitat; function; seagrass; waves; attenuation, 565
HART, B., *see* SHILLABEER, N., 105
Hawaii; mixing model; anthropogenic; nutrients; groundwater, 409
HAWKINS, S. J., *see* CONLAN, K., 435
HIRCHE, H.-J., Egg production of *Eurytemora affinis*—effect of K-strategy, 395
HOPKINSON, C. S., *see* NEWELL, S. Y., 127
HOWES, B. L., *see* ORSON, R. A., 453
Hydrography; freshets; natural perturbations; Gernika estuary; plankton dynamics, 425
IMBERGER, J., *see* ALLANSON, B. R., 253
Induced laminar flow; Reynolds number; tube aspect ratio; boundary layer; suspensoids; thalassinidea; burrows; Bernoulli effect, 253
Instruments; electromagnetics; erosion; siltation, 353
Internal waves; estuaries; benthic boundary layer; bed friction; wave-current interactions, 17
Intertidal; southern Africa; recruitment; growth; barnacles, 167
Intertidal environment; deposition; removal of contaminants; mudland sandflats; microbial degradation; trace elements, 371
ISHIWATARI, R., *see* TAKADA, H., 141
Jiangsu Province, China; tidal mud flat; tidal current; sediment concentration; residual circulation, 225
JICKELLS, T. D., *see* FICHEZ, R., 577
JOHNS, B., RAO, A. D. and RAO, G. S., On the occurrence of upwelling along the east coast of India, 75
KERNER, M. and WALLMANN, K., Remobilization events involving Cd and Zn from intertidal flat sediments in the Elbe estuary during the tidal cycle, 371
KERR, J. D., *see* BLABER, S. J. M., 605
KIM, S. C., *see* WRIGHT, L. D., 17
Kinetics; estuaries; England coast; trace metals, 37
KJERFVE, B., *see* ALTHAUSEN, J. D., 517
KJERFVE, B., *see* PILLAY, S., 331
Kriging; spatial distribution; *Munida intermedia*; *Munida sarsi*; Galician continental shelf; NW Spain; geostatistical analysis; variograms, 637
Lagoons; pollution; plankton; England coast; estuaries; docks, 435
Land-derived pollutants; synthetic detergents; surfactants; organic micropollutants; Tokyo Bay; linear alkylbenzenes (LABs); linear alkylbenzenesulphonates (LAS), 141
Larval transport; coastal lagoon; bivalves; *Ruditapes*; Mediterranean; recruitment, 289
Laser analysis; dry-sieving, comparison; sediment; particle size, 105
LASSERRE, G., *see* BACH, P., 55
LAW, C. S., REES, A. P. and OWENS, N. J. P., Nitrous oxide: estuarine sources and atmospheric flux, 301

LAYBOURN-PARRY, J., ROGERSON, A. and CRAWFORD, D. W., Temporal patterns of protozooplankton abundance in the Clyde and Loch Striven, 533
LEBO, M. E. and SHARP, J. H., Modeling phosphorus cycling in a-well-mixed coastal plain estuary, 235
LEGENDRE, P., *see* BACH, P., 55
Linear alkylbenzenes (LABs); linear alkylbenzenesulphonates (LAS); land-derived pollutants; synthetic detergents; surfactants; organic micropollutants; Tokyo Bay, 141
Linear alkylbenzenesulphonates (LAS); land-derived pollutants; synthetic detergents; surfactants; organic micropollutants; Tokyo Bay; linear alkylbenzenes (LABs), 141
LLANSÓ, R. J., Effects of hypoxia on estuarine benthos: the lower Rappahannock River (Chesapeake Bay), a case study, 491
Loch Striven; Clyde; Scotland coast; estuarine; flagellates; ciliates; dinoflagellates; flocs, 233
Long Term Ecological Research (LTER); North Inlet, SC; Bass Creek, SC; suspended sediment concentration; sediment transport rate; tidal estuary; *Spartina* marsh; errors, 331
MCCARTHY, R. K., *see* GARVINE, R. W., 157
Macrobenthos; population dynamics; burrowing behaviour; Chesapeake Bay, U.S.A. coast; dissolved oxygen, 491
Macrobenthos; seagrass; community; drilling mud; Santa Rosa Sound, Florida, 315
MARTÍN, D., *see* PALACÍN, C., 1
Mass balance; regeneration; Delaware Bay; phosphorus; models; estuaries, 235
Mediterranean; recruitment; larval transport; coastal lagoon; bivalves; *Ruditapes*, 289
Mediterranean coast; nematoda; meiobenthos; eutrophication; population; density and variations, 1
MEHTA, A., *see* WOLANSKI, E., 473
Meiobenthos; eutrophication; population; density and variations; Mediterranean coast; nematoda, 1
Microbial degradation; trace elements; intertidal environment; deposition; removal of contaminants; mudland sandflats, 371
Microdetritus; stratification; advection; Adriatic Sea; eutrophication; phytoplankton; chlorophyll *a* distribution, 191
MILLET, B., *see* BORSA, P., 289
MILLWARD, G. E., GLEGG, G. A. and MORRIS, A. W., Zn and Cu removal kinetics in estuarine waters, 37
MIQUEL, J.-C., *see* PUŠKARIĆ, S., 267
Mixing model; anthropogenic; nutrients; groundwater; Hawaii, 409
Models; estuaries; mass balance; regeneration; Delaware Bay; phosphorus, 235
Models; estuarine circulation; tides, 113
MORRIS, A. W., *see* MILLWARD, G. E., 37
Mudland sandflats; microbial degradation; trace elements; intertidal environment; deposition; removal of contaminants, 371
Mud suspension; Australia coast; dredging; sea dumping; settling; sedimentation, 473
Munida intermedia; *Munida sarsi*; Galician continental shelf; NW Spain; geostatistical analysis; variograms; kriging; spatial distribution, 637
Munida sarsi; Galician continental shelf; NW Spain; geostatistical analysis; variograms; kriging; spatial distribution; *Munida intermedia*, 637
NARAYANA, A. C., *see* PANDARINATH, K., 363
Natural perturbations; Gernika estuary; plankton dynamics; hydrography; freshets, 425
NEIRA, F. J. and POTTER, I. C., Movement of larval fishes through the entrance channels of a seasonally open estuary in Western Australia, 213
Nematoda; meiobenthos; eutrophication; population; density and variations; Mediterranean coast, 1
New England Coast; paleoecology; geomorphology; plant community structure, 453
NEWELL, S. Y., HOPKINSON, C. S. and SCOTT, L. A., Patterns of nitrogenase activity (acetylene reduction) associated with standing, decaying shoots of *Spartina alterniflora*, 127
Nitrification; denitrification; estuarine sediment; atmospheric flux; Tamar estuary; South West England; nitrous oxide; nutrients, 301
Nitrogen fixation; acetylene reduction; saltmarsh; *Spartina alterniflora*; decomposition; clay deposition, 127

Nitrous oxide; nutrients; nitrification; denitrification; estuarine sediment; atmospheric flux; Tamar estuary; South West England, 301
Northern Adriatic Sea; annual cycle; particulate flux, 267
North Inlet, SC; Bass Creek, SC; suspended sediment concentration; sediment transport rate; tidal estuary; *Spartina* marsh; errors; Long Term Ecological Research (LTER), 331
Numerical model; coastal upwelling; estuarial discharge; Bay of Bengal, 75
Nursery areas; Zambezi River; penaeid shrimp; river regulation; salinity, 91
Nutrient cycling; estuary; algal bloom, 577
Nutrients; groundwater; Hawaii; mixing model; anthropogenic, 409
Nutrients; nitrification; denitrification; estuarine sediment; atmospheric flux; Tamar estuary; South West England; nitrous oxide, 301
ODEBRECHT, C., *see* ABREU, P. C., 621
OGURA, N., *see* TAKADA, H., 141
OLASO, I., *see* FREIRE, J., 637
Organic matter; calcium; Gangolli estuary; clay minerals; trace elements, 363
Organic micropollutants; Tokyo Bay; linear alkylbenzenes (LABs); linear alkylbenzenesulphonates (LAS); land-derived pollutants; synthetic detergents; surfactants, 141
ORIVE, E., *see* DE MADARIAGA, I., 425
ORSON, R. A. and HOWES, B. L., Salt marsh development studies at Waquoit Bay, Massachusetts: influence of geomorphology on long-term plant community structure, 453
OWENS, N. J. P., *see* LAW, C. S., 301
Oyster fishery; freshwater inflows; Apalachicola Bay, 179
PALACÍN, C., GILI, J.-M. and MARTÍN, D., Evidence for coincidence of meiofauna spatial heterogeneity with eutrophication processes in a shallow-water Mediterranean bay, 1
Paleoecology; geomorphology; plant community structure; New England Coast, 453
Pandalidae; *Parapandalus narval*; Greece coast; fecundity; reproductive effort; depth, 593
PANDARINATH, K. and NARAYANA, A. C., Clay minerals and trace metal association in the Gangolli estuarine sediments, west coast of India, 363
Parapandalus narval; Greece coast; fecundity; reproductive effort; depth; Pandalidae, 593
Particle size; laser analysis; dry-sieving, comparison; sediment, 105
Particulate flux; northern Adriatic Sea; annual cycle, 267
Particulate organic carbon; estuaries; Brazil coast; South American South coast; bacteria; phytoplankton; protozoa; seston, 621
Path analysis; eel; fishing gear; fishing effort; fishing strategy; Thau lagoon; contingency periodogram, 55
Penaeid shrimp; river regulation; salinity; nursery areas; Zambezi River, 91
Phosphorus; models; estuaries; mass balance; regeneration; Delaware Bay, 235
Phytoplankton; chlorophyll *a* distribution; microdetritus; stratification; advection; Adriatic Sea; eutrophication, 191
Phytoplankton; protozoa; seston; particulate organic carbon; estuaries; Brazil coast; South American South coast; bacteria, 621
PILLAY, S., GARDNER, L. R. and KJERFVE, B., The effect of cross-sectional velocity and concentration variations on suspended sediment transport rates in tidal creeks, 331
Plankton dynamics; hydrography; freshets; natural perturbations; Gernika estuary, 425
Plankton; England coast; estuaries; docks; lagoons; pollution, 435
Plant community structure; New England Coast; peleoecology; geomorphology, 453
Pollution; plankton; England coast; estuaries; docks; lagoons, 435
Population dynamics; burrowing behaviour; Chesapeake Bay, U.S.A. coast; dissolved oxygen; macrobenthos, 491
Population; density and variations; Mediterranean coast; nematoda; meiobenthos; eutrophication, 1
POTTER, I. C., *see* NEIRA, F. J., 213
Protozoa; seston; particulate organic carbon; estuaries; Brazil coast; South American South coast; bacteria; phytoplankton, 621
PUŠKARIĆ, S., FOWLER, S. W. and MIQUEL, J.-C., Temporal changes in particulate flux in the northern Adriatic Sea, 267
RAO, A. D., *see* JOHNS, B., 75

RAO, G. S., *see* JOHNS, B., 75
Recruitment; growth; barnacles; intertidal; southern Africa, 167
Recruitment; larval transport; coastal lagoon; bivalves; *Ruditapes*; Mediterranean, 289
REES, A. P., *see* LAW, C. S., 301
Regeneration; Delaware Bay; phosphorus; models; estuaries; mass balance, 235
Removal of contaminants; mudland sandflats; microbial degradation; trace elements; intertidal environment; deposition, 371
RENSHUN, Z., Suspended sediment transport processes on tidal mud flat in Jiangsu Province, China, 225
Reproductive effort; depth; Pandalidae; *Parapandalus narval*; Greece coast; fecundity, 593
Reproductive strategy; *Eurytemora*; egg production, 395
Residual circulation; Jiangsu Province, China; tidal mud flat; tidal current; sediment concentration, 225
REVELANTE, N. and GILMARTIN, M., The lateral advection of particulate organic matter from the Po Delta region during summer stratification, and its implications for the northern Adriatic, 191
Reynolds number; tube aspect ratio; boundary layer; suspensoids; thalassinidea; burrows; Bernoulli effect; induced laminar flow, 253
RIDD, P., *see* WOLANSKI, E., 473
RIDD, P. V., A sediment level sensor for erosion and siltation detection, 353
RIDDLE, A. M., *see* SHILLABEER, N., 105
River flow; diffusivity; effective longitudinal; Delaware Bay; salinity variations; estuaries, 157
River regulation; salinity; nursery areas; Zambezi River; penaeid shrimp, 91
ROGERSON, A., *see* LAYBOURN-PARRY, J., 533
Ruditapes; Mediterranean; recruitment; larval transport; coastal lagoon; bivalves, 289
SALINI, J. P., *see* BLABER, S. J. M., 605
Salinity; nursery areas; Zambezi River; penaeid shrimp, river regulation, 91
Salinity; temperature; fish distribution; estuarine habitat; seasonal patterns; turbidity, 545
Salinity variations; estuaries; river flow; diffusivity; effective longitudinal; Delaware Bay, 157
Saltmarsh; *Spartina alterniflora*; decomposition; clay deposition; nitrogen fixation; acetylene reduction, 127
Santa Rosa Sound, Florida; macrobenthos; seagrass; community; drilling mud, 315
Scotland coast; estuarine; flagellates; ciliates; dinoflagellates; flocs; Loch Striven; Clyde, 533
SCOTT, L. A., *see* NEWELL, S. Y., 127
Sea dumping; settling; sedimentation; mud suspension; Australia coast; dredging, 473
Seagrass; community; drilling mud; Santa Rosa Sound, Florida; macrobenthos, 315
Seagrass, tropical; fish; species composition; biomass; Gulf of Carpentaria; Australia coast, 605
Seagrass; waves; attenuation; habitat; function, 565
Seasonal patterns; turbidity; salinity; temperature; fish distribution; estuarine habitat, 545
Sediment; particle size; laser analysis; dry-sieving, comparison, 105
Sediment concentration; residual circulation; Jiangsu Province, China; tidal mud flat; tidal current, 225
Sediment transport rate; tidal estuary; *Spartina* marsh; errors; Long Term Ecological Research (LTER); North Inlet, SC; Bass Creek, SC; suspended sediment concentration, 331
Sedimentation; mud suspension; Australia coast; dredging; sea dumping; settling, 473
Seston; particulate organic carbon; estuaries; Brazil coast; South America South coast; bacteria; phytoplankton; protozoa, 621
Settling; sedimentation; mud suspension; Australia coast; dredging; sea dumping, 473
SHARP, J. H., *see* LEBO, M. E., 235
SHETYE, S. R. and GOUVEIA, A. D., On the role of geometry of cross-section in generating flood-dominance in shallow estuaries, 113
SHILLABEER, N., HART, B. and RIDDLE, A. M., The use of a mathematical model to compare particle size data derived by dry-sieving and laser analysis, 105
Siltation; instruments; electromagnetics; erosion, 353
SKINNER, D., *see* ALLANSON, B. R., 253
Slack water; suspended sediment; tides; transmissivity; turbidity maximum zone; U.S.A. coast; fresh water discharge; partially mixed estuary, 517

South America South coast; bacteria; phytoplankton; protozoa; seston; particulate organic carbon; estuaries; Brazil coast, 621
South West England; nitrous oxide; nutrients; nitrification; denitrification; estuarine sediment; atmospheric flux; Tamar estuary, 301
Southern Africa; recruitment; growth; barnacles; intertidal, 167
Spain, NW; geostatistical analysis; variograms; kriging; spatial distribution; *Munida intermedia*; *Munida sarsi*; Galician continental shelf, 637
Spartina marsh; errors; Long Term Ecological Research (LTER); North Inlet, SC; Bass Creek, SC; suspended sediment concentration; sediment transport rate; tidal estuary, 331
Spartina alterniflora; decomposition; clay deposition; nitrogen fixation; acetylene reduction; saltmarsh, 127
Spatial distribution; *Munida intermedia*; *Munida sarsi*; Galician continental shelf; NW Spain; geostatistical analysis; variograms; kriging, 637
Species composition; biomass; Gulf of Carpentaria; Australia coast; tropical seagrass; fish, 605
Stratification; advection; Adriatic Sea; eutrophication; phytoplankton; chlorophyll *a* distribution; microdetritus, 191
Surface waves; wind stress; wind-driven currents, 347
Surfactants; organic micropollutants; Tokyo Bay; linear alkylbenzenes (LABs); linear alkylbenzenesulphonates (LAS); land-derived pollutants; synthetic detergents, 141
Suspended sediment concentration; sediment transport rate; tidal estuary; *Spartina* marsh; Long Term Ecological Research (LTER); North Inlet, SC; Bass Creek, SC, 331
Suspended sediment; tides; transmissivity; turbidity maximum zone; U.S.A. coast; fresh water discharges; partially mixed estuary; slack water, 517
Suspensoids; thalassinidea; burrows; Bernoulli effect; induced laminar flow; Reynolds number; tube aspect ratio; boundary layer, 253
Synthetic detergents; surfactants; organic micropollutants; Tokyo Bay; linear alkylbenzenes (LABs); linear alkylbenzenesulphonates (LAS); land-derived pollutants, 141
TAKADA, H., ISHIWATARI, R. and OGURA, N., Distribution of linear alkylbenzenes (LABs) and linear alkylbenzenesulphonates (LAS) in Tokyo Bay sediments, 141
Tamar estuary; South West England; nitrous oxide; nutrients; nitrification; denitrification; estuarine sediment; atmospheric flux, 301
Temperature; fish distribution; estuarine habitat; seasonal patterns; turbidity; salinity, 545
Thalassinidea; burrows; Bernoulli effect; induced laminar flow; Reynolds number; tube aspect ratio; boundary layer; suspensoids, 253
Thau lagoon; contingency periodogram; path analysis; eel; fishing gear; fishing effort; fishing strategy, 55
THESSALOU-LEGAKI, M., Reproductive variability of *Parapandalus narval* (Crustacea: Decapoda) along a depth gradient, 593
Tidal currents; sediment concentration; residual circulation; Jiangsu Province, China; tidal mud flat, 225
Tidal estuary; *Spartina* marsh; errors; Long Term Ecological Research (LTER); North Inlet, SC; Bass Creek, SC; suspended sediment concentration; sediment transport rate, 331
Tidal mud flat; tidal current; sediment concentration; residual circulation; Jiangsu Province, China, 225
Tides; models; estuarine circulation, 113
Tides; transmissivity; turbidity maximum zone; U.S.A. coast; fresh water discharge; partially mixed estuary; slack water; suspended sediment, 517
Tokyo Bay; linear alkylbenzenes (LABs); linear alkylbenzenesulphonates (LAS); land-derived pollutants; synthetic detergents; surfactants; organic micropollutants, 141
Trace elements; intertidal environment; deposition; removal of contaminants; mudland sandflats; microbial degradation, 371
Trace elements; organic matter; calcium; Gangolli estuary; clay minerals, 363
Trace metals; kinetics; estuaries; England coast, 37
Transmissivity; turbidity maximum zone; U.S.A. coast; fresh water discharge; partially mixed estuary; slack water; suspended sediments; tides, 517
Transport; Western Australian coast; estuaries; fish larvae, 213